NHS Kent and Medway

Low Molecular Weight Prescribing Pathways in East Kent

Obstetric LMWH Prescribing Pathways

The pathways in the table below were agreed at a meeting on 23.6.22 with K&M LMC Representative, Consultant Haematologist EKHUFT, GP K&M CCG Clinical Prescribing Lead, O&G Consultant EKHUFT, Head of Quality Maternity and Neonatal Services Kent and Medway Local Maternity System (LMNS), Clinical lead for Kent and Medway LMS, Senior Programme Manager Maternity K&M ICB, Neonatal and Perinatal Mental Health, and O&G Care Groups Operations Director EKHUFT.

LMWH Obstetric Rxing Pathway	Prescribing Pathway agreed 23.6.22
Obstetrics - high risk from day 1 of pregnancy including IVF patients (LMWH during IVF	All prescribing in East Kent to remain with EKHUFT
treatment will be prescribed by the IVF centre)	
Obstetrics -High risk identified at 8-10 week midwife booking appointment	All prescribing in East Kent to remain with EKHUFT
Obstetrics - treatment of DVT/PE	All prescribing in East Kent to remain with EKHUFT
Obstetrics - post natal treatment	No change- full supply given by EKHUFT
Obstetrics- COVID	No change- full supply given by EKHUFT

If an un-booked patient who needs thromboprophylaxis from day 1 (pathway 1) contacts a GP, the GP may contact Advice and Guidance (Haematology) at EKHUFT.



Non-obstetric LMWH Prescribing Pathways

The following pathways were agreed by the former East Kent Prescribing Group (EKPG) in January 2020. An agreement with Oncology for patients undergoing chemotherapy was not reached, and the EKPG therefore agreed an interim solution.

INDICATION	INITIATION BY	AMOUNT TO BE SUPPLIED ON INITIATION	REPEAT SUPPLIES BY	COMMENTS
	GP prior to ultra sound scan	Until diagnosis/ therapeutic		
Suspected DVT treatment	on advice of DVT clinic	INR	Not required	
	EKHUFT	Until diagnosis/ therapeutic		
Suspected PE treatment		INR	Not required	
	EKHUFT	2-6 week supply as		
Surgical procedures -		appropriate to surgical		
prophylaxis		procedure	Not required	
Surgical procedures -	EKHUFT			
Bridging and peri-operative				
use		As appropriate	Not required	
	Interim arrangement. East Kent Prescribing Group advise that while it is			
	preferable that supplies are made by EKHUFT, GPs could prescribe if they have received written information confirming the			
Oncology - For patients	drug, dose and monitoring			
undergoing chemotherapy	needed.			
Oncology - Patient requiring	GP - on receipt of letter			
ongoing enoxaparin	from haematologist			
treatment following	confirming need for ongoing			
	treatment and duration	On- going treatment	GP	

Approved by: IMOC Approval date: Feb 2024 Review date: Feb 2026



Kent and Medway

completion of				
chemotherapy				
Contraindications to oral	EKHUFT			For existing patients consider review by haematology as DOAC may
		2 wooles	CD	
anticoagulants		2 weeks	GP	now be appropriate
	EKHUFT; KCHFT; Other			
Lower limb immobilisation	Minor Injuries providers	Full supply	n/a	
Superficial	EKHUFT			
Thrombophlebitis		Full supply	n/a	

Approved by: IMOC Approval date: Feb 2024 Review date: Feb 2026